

ENT SURGICAL CONSULTANTS

Thomas K. Kron, MD, FACS
 Michael G. Gartlan, MD, FAAP, FACS
 Rajeev H. Mehta, MD, FACS
 Scott W. DiVenere, MD
 Sung J. Chung, MD
 Ankit M. Patel, MD

2201 Glenwood Ave., Joliet, IL 60435
 (815) 725-1191, (815) 725-1248 fax

1300 Copperfield Ave., Suite 3060, Joliet, IL 60432
 (815) 727-6031

119 E. Jefferson St., Morris, IL 60450
 (815) 941-1972

ORAL ANTIHISTAMINES (11/08)

General Information

- Can be used on an as needed basis, yet best if used prior to significant allergy exposure (i.e., cats, cutting grass, picnics, etc).
- Oral antihistamines help relieve symptoms of sneezing, clear nasal discharge, itchy/watery/red eyes, and itchy nose/palate/and/or throat, yet *have no effect on nasal congestion*.
- Some antihistamines are combined with a decongestant ("D") to help with allergy related nasal congestion, yet this increases the viscosity of the mucus and may lead to a higher incidence of sinusitis or nasal infection.
- Use only one antihistamine at a time.
- Oral antihistamines are only useful for true allergy symptoms. These medications are not indicated and are not appropriate for use in colds or upper respiratory infections.

First Generation Sedating Oral Antihistamines

- Many are available over-the-counter (OTC) without a prescription (Benadryl, Tavist, etc).
- First generation antihistamines often cause drowsiness, sedation, reduced motivation, as well as impaired motor and cognitive function. This poses significant safety hazards while driving or operating machinery.
- Become less effective if used over a prolonged period of time due to tolerance.
- Slow down the normal clearance of mucus from the nose and sinuses and can make you more susceptible to chronic sinusitis.

Second Generation Non-Sedating Oral Antihistamines

Brand Name	Dose & Forms	Indicated Age (yr)	Recommended Dose	Sedation	Available OTC
Claritin®, or generic loratidine	1 mg/mL syrup	2 to 5	2.5-5 mg once daily	No	Yes
	10 mg tablet	≥ 6	10 mg once daily		
	10 mg RediTabs®				
Claritin-D® 12 Hour	5 mg/120 mg extended-release tablet	≥ 12	One tablet twice daily	No	Yes
Claritin-D® 24 Hour	10 mg/240 mg extended-release tablet	≥ 12	One tablet daily	No	Yes
Zyrtec®, or generic fexofenidine	5 mg/5 ml syrup	2 to 5	2.5 mg to 5 mg once daily (or 2.5 mg every 12 hours)	Mild	Yes
	5 mg tablets	6 to 12	1-2 tablets daily		
	10 mg tablets	≥ 12	One tablet daily		
Zyrtec D® 12 Hour	5 mg/120 mg extended release tablet	≥ 12	One tablet twice daily	Mild	Yes
Clarinex	2.5 mg/ 5 mL syrup	6 mo-11 yo	1.0-2.5 mg (2 -5 mL) once daily	No	No
	5 mg tablet	>_12	One 5 mg tablet once daily		
	5 mg RediTabs®				
Allegra®	30 mg tablet	6 to 11	30 mg twice daily	No	No
	60 mg capsule or tablet	≥ 12	60 mg twice daily		
	180 mg tablet	≥ 12	180 mg once daily (SAR only)		
Allegra-D®	60 mg/120 mg extended-release tablet	≥ 12	One tablet twice daily	No	No
Xyzal	5 mg tablet	6 mo-11 yo	2.5 mg once daily	No	No
		>_12	5 mg once daily		

Contraindications

- Due to the decongestant component ("D") of some of the above preparations, this medication is contraindicated in patients with urinary retention, narrow-angle glaucoma, and concurrent use of monoamine oxidase (MAO) inhibitor antidepressants.